


[The UMBS Adaptive Aspen Management Experiment \(AAME\)](#) [updated 2/10/20]

By Luke Nave, with design assistance from Stephen Handler, Inés Ibañez, Maria Janowiak, Knute Nadelhoffer, Dean Reid, Adam Schubel, Karie Slavik, Chris Swanston, Jason Tallant

Summary: This document describes forest manipulations on the interlobate moraine, known locally as the Pellston Hill, which were implemented in 2019. Goals of the manipulations are many and complementary. First, make UMBS forests less vulnerable to climate change (ongoing and future) by implementing the Northern Institute of Applied Climate Science (NIACS) Climate Change Response Framework (CCRF). The CCRF is based upon forest adaptation concepts of *resistance*, *resilience*, and *transition* (Figure 10 in Swanston et al. 2016). Testing these concepts with experimental manipulations, while maintaining untreated *reference* areas, makes [UMBS a formal demonstration site](#) in the NIACS network, showcasing alternatives for aspen management in the face of climate change. The second goal is to generate science products pertaining to harvest impacts on hydrology and biogeochemistry using revenue gained from timber sales. Third, create research opportunities for classes, students, and investigators, and fourth, build relationships with partners in the forestry industry and forest-climate science and outreach institutions that will serve UMBS interests and advance climate-adaptive forest management. Below, I move from general to specific in describing the areas, objectives, and research questions associated with the treatments.

Management Area: Figure 1 shows the location of the aspen-dominated 183 ha management area, subdivided into 18 units ranging in size from ~2 – 73 ha each. Unit shapes and sizes are constrained by existing stand ages, research sites, and questions. Seven units are young stands regenerated after clearcuts in the early 1970s (18, 19) or early 1980s (units 2, 4, 6, 8, 10, 12). The young units total ~25 ha and provide a second structure / age class in the current landscape mosaic, which is dominated by mature aspen dating to the 1908-1919 period (Art Cooper plot data). Of these areas of old aspen,



presently totaling ~157 ha in the management area, 30 ha were harvested during the first set of manipulations. Harvested units are colored in pink (3, 7, 9, 13), while reference units are dark green (1, 5, 11, 14, 15, 16, 17). Areas are allocated into cut (pink) vs. reference (dark green) units according to several criteria. First, ensure that research locations, such as long-term plots (orange and purple points including Landscape Ecosystems, 50 Year and Art Cooper Plots) and study areas (e.g.,

Honeysuckle Creek headwaters and Burt Barnes' aspen clones in units 14 and 15, respectively) are not affected by manipulations. Second, protect unique hydrologic features, such as Surprise Pond (in unit 11) and a swamp / ephemeral stream between units 6 and 7. Third, allocate treatments across space in such a way that the distinct forest adaptation *concepts* (resistance, resilience, transition, reference) can be tested at the landscape level, by implementing specific *strategies* and *approaches* at the unit level, while allowing for stand-level (within-unit) flexibility in *tactics* (again see Swanston et al. 2016). This experimental framework allows *strategies* and *approaches* to be evaluated across a range of ecosystem conditions within each unit (e.g., tree composition, soils, hydrology), while affording *tactical* flexibility for the consulting forester and logging operators to respond to on-the-ground constraints. Fourth, allocate sufficient overall area to treatments as to make the necessary timber sale economically attractive to the operators who will bid on and perform the work. The details below describe a management plan that optimizes across the above four criteria.

Treatments: The overall long-term goal of harvesting is to modify the existing, aspen-dominated forest of the management area into a matrix of conditions better positioned to maintain their ecological functions as climate continues to change. The more immediate goal of harvesting is to provide locations and funds for studying forest management impacts on carbon forms, turnover, and mobility in soils and groundwater. Treatments will produce immediate changes in forest composition, structure, hydrology, and biogeochemistry that then play out over longer time frames, as the units undergo succession. In 2019, the treatments removed standing biomass (mostly aspen) in order to either: 1) regenerate young, homogenous-structure aspen/birch cover types (*resistance*); 2) regenerate aspen-dominated cover, but with greater structural and compositional diversity (*resilience*); 3) encourage succession to more structurally and compositionally diverse cover types (*transition*). Specific treatments varied according to the cover types, soils, and hydrology present on the units, which are represented well by the landscape-level ecosystem types (Figure 2) in the management area. For a thorough description of ecosystem types, refer to Pearsall (1995); for consideration of their soils and hydrology (Hofmeister et al. 2019a), and groundwater chemistry (Hofmeister et al. 2019b); for their application to forest growth rates Nave et al. (2017). Briefly, key ecosystem factors in the management area include: 1) landform, including parent material, slope and aspect; 2) hydrology; 3) soil texture. These three factors affect current vegetation cover, and will continue to do so into the future regardless of treatment. Regarding landform, the key difference in the management area is between moraine vs. outwash, dune, and beach ridge landforms. Moraine ecosystems (types 109, 113, 116, 118 in Figure 2) have nutrient-rich, fine-textured glacial till near the surface; these are level to gently sloping and because slopes are generally N- and E-facing, the topoclimate is cooler and moister. The presence of a seasonal (109, 113) to permanent (116, 118) water table ≤ 2 m of the surface is closely tied to the depth to till, which in some places is present as a fragipan or otherwise dense enough to exclude deep rooting. Bigtooth aspen is very productive on these ecosystems, and accessory species range from mesic northern hardwoods (beech, maples, red oak, basswood) to riparian and wetland species (trembling aspen, green and black ash, balsam fir). Outwash (36, 39, 47, 59, 60), dune (74), and beach ridge (75, 76) ecosystems have their differences, but share a common difference from moraine ecosystems in that they all have deep, sandy parent materials. Outwash landforms are mostly nearly level; dunes and beach ridges have locally steep slopes of any aspect; overall all of these ecosystems are considerably drier and lower in fertility than


moraine ecosystems and typically support lower-stature and lower-density cover types dominated by bigtooth aspen, red maple and oak, and white and red pines. A seasonal (spring melt) water table exists in these ecosystems only in sporadic locations directly adjacent to (topographically below) moraine ecosystems. Following this brief overview of ecosystem types, some short definitions of the forest adaptation concepts that motivated the more detailed, unit-level management treatments are in order:

Resistance actions improve the defenses of an ecosystem against expected changes, in order to maintain relatively unchanged conditions. In this context, *resistance* means to resist climate-driven changes in forest composition by regenerating a vulnerable cover type in a protected location.

Resilience actions allow some change, but help an ecosystem return to its prior state after a disturbance, either naturally or through management. In this context, *resilience* means to maintain aspen-dominated stands as such, but to increase their structural and compositional diversity so their functions (e.g., C uptake, hydrology) can better recover from future climate change impacts and other disturbances.

Transition actions actively facilitate change, in order to help ecosystems adapt to changing conditions. In this context, *transition* means to convert aspen-dominated stands more diverse cover types that will have a higher probability of maintaining ecosystem functions despite future climate and disturbances.

Reference describes a state of no intervention, in which ecosystems undergo climate change and other disturbances and their responses to those drivers are monitored.



Unit 3 Concept: *Resilience*. This 6 ha unit was the least diverse, in terms of ecosystem types (3) and pre-harvest forest cover, which was overwhelmingly dominated by very tall, large-diameter bigtooth aspen. Red oak, beech, and white birch were present as scattered codominants, and understory (pole and sapling) species primarily included red maple, beech, and striped maple. Most of the unit is level to gently sloping, with a general E-facing aspect, but with some locally very steep (>25%) slopes in the northern portion of the unit. All portions are on the moraine. The ecosystem types present (59, 60, and 109) are excessively- to well-drained, and except in the ravine in the northern portion (and then only in

spring) there is likely no water table present. Thus, given the relative lack of species and structural diversity, and the relatively dry soil/ecosystem types present, this unit is particularly vulnerable to climate change. The goal then was to remove most of the over-mature aspens, which are increasingly being blown over due to their height and the restrictive subsoil (a fragipan is present in some areas of this unit), resulting in a more spatially complex forest structure and species composition that will be better positioned to recover from climate change-induced disturbances (now and in the future). The treatment: 1) removed all diseased beech and 2) ~75% of the aspen on the overall unit. Aspen removals were most intensive at the western edge of the unit, with an increasing density of residual large aspens as one moves eastward through the unit. This feathering effect (eastward-increasing residual aspen density) was intended to create a more diffuse boundary on the unit's eastern edge than if all aspen were removed from the entire unit, in order to increase boundary layer resistance and make the units located to the east (16 and 9) less vulnerable to strong winds. Overall aspen retention was ~19 trees per hectare. On unit 3, stems of all other co-dominant and understory species were retained to the degree possible; these include primarily red oak and red maple, but also some non-diseased beech (~15 per ha). Future work may involve the planting of southern genotypes of species already present at UMBS (e.g., red oak, white pine, bigtooth aspen, sugar maple), and introductions of species that reach range limits near to the south (e.g., white oak, Hill's oak, shagbark hickory) or farther afield. Planted trees will need to be protected from browse; this may be accomplished by planting within any of 38 slash piles (each 6-10 m diameter and 1-2 m tall, all have been GPS'd) that were created across the unit during harvesting. Cutting on this unit was completed in March-April 2019.

Unit 9 Concept: *Resilience*. This 7 ha unit has slightly higher diversity of ecosystem (4) and cover types than Unit 3. It occupies level to gently sloping, mostly N/E facing lower flanks of the moraine. There are some local areas of S-facing slopes and outwash channels in the northeastern portion of the unit, where the edge of the moraine meets outwash and dune ecosystems. Before harvesting, bigtooth aspen dominated throughout. Aspen was very tall and large-diameter on the moraine (ecosystem 109), where red oak and beech were codominant, and red maple, beech, and striped maple formed the understory. On the drier outwash (39), dune (74), and deep outwash over till ecosystems (59), aspen density, height, and diameter was lower; codominant and understory taxa were red maple, red oak, white pine, and white birch. Given the slightly higher level of ecosystem and cover type diversity here, it is likely more resilient (at the unit level) to climate change and its impacts than Unit 3, but still vulnerable due to the dominance of bigtooth aspen throughout. Further, its topographic variability (both N- and S-facing slopes) may afford opportunities to test the performance of different planted species. Similar to Unit 3, treatment of this unit was intended to increase spatial, compositional, and structural diversity, though the harvest prescription differed somewhat. The prescription was to: 1) remove ~95% of aspen, with a more or less uniform distribution of retained aspen (~5 per ha). 2) Remove all diseased beech [retaining ~12 non-diseased beech per ha]. 3) In drier eastern portions of the unit, remove ~20% of pole-sized red and white pine in densely stocked areas totaling 1-3 ha, in order to release residual pines and oaks from shading; also remove ~15 red oaks from this area to provide firewood for the UMBS campus and a small quantity of high-value sawlogs. Cutting on this unit was completed in April-May 2019.

Unit 7 Concept: *Transition*. This 7 ha unit has the second-highest level of within-unit ecosystem diversity, with diverse cover types and locally complex topography. The majority of the unit is on level to gently sloping N/E facing lower moraine slopes (ecosystem 109), where very tall, large-diameter bigtooth aspen dominated but with a notable presence of sapling, pole, and even co-dominant stems of red oak, red maple, yellow and paper birch, ironwood, striped maple, basswood, and hemlock. An eastward-running meltwater channel that is adjacent to (but outside) the unit's northern edge holds nearly perennially saturated soils, and supports a cover of trembling aspen, balsam fir, ash (green and black), red maple, white pine, hemlock, and white cedar. The eastern 1/3 of the unit includes gentle E, S, and W-facing slopes that flank a low-lying, northward-running (dry) outwash channel, bounded along its S and E sides by old dunes. In these areas of sandier parent materials the stem density and canopy cover was lower, and dominated by bigtooth aspen. The aspen was of lower stature and in some areas was codominant with red pine and even a few jack pines (apparently plantation), while red oak, white pine, red maple and balsam fir were accessory taxa. Given the diversity of existing cover types, topographic and hydrologic positions, and the presence of considerable accessory species diversity, this unit lent itself to management intended to transition away from dominance by bigtooth aspen, towards more diverse cover types. In order to achieve this transition, removals on this unit were exclusively targeted at aspen (all of it over-mature and some of it beginning to break up or fall over) and diseased beech. All other stems of all species, from co-dominant to understory sizes, were retained to the degree possible, except in the northeastern portion of the unit, where some localized areas of red pine (<2 ha) were lightly thinned to release over-crowded residual stems from light competition. In this part of the unit, care was taken to leave a dense overstory of red and white pines in locations that shade the cool, wet hydric ecosystems between this unit and Unit 6. Cutting on this unit was completed in March 2019.

Unit 13 Concept: *Resistance*. This 10 ha unit has the highest level of within-unit ecosystem diversity (9 ecosystem types), and accordingly, much within-unit variety in soils, hydrology, and vegetation. It also has a favorable physiographic position (NE-facing aspect with locally steep slopes and high water tables), making it well suited to attempts to resist climate change-driven drought and heat. Further, the unit is within part of the Honeysuckle Creek Watershed (HCW) where we have sampled soils and groundwater since 2014. It is isolated from the HC headwaters by a moraine/dune ridge to its south, creating a physically separate sub-basin where the impacts of clearcutting on hydrology, soil C turnover and mobilization are now being studied in real time, and compared to adjacent stands (reference and 1980s cuts) on ecosystem types that extend into adjacent units. In this case, all of these factors align well with a "business-as-usual," resistance-oriented management strategy. In other words, nearly all merchantable stems were cut and removed from this unit, with the intent of regenerating a mixed aspen-birch cover type. This cover type is generally viewed as a climate change loser in northern Lower Michigan (Handler et al. 2014). We will test that prediction with a uniform prescription, applied across the range of ecosystems presented in the unit (including drier outwash and dune ecosystems on the unit's E edge), intending to learn whether this approach to aspen-birch management can sustain it despite ongoing climate change. The prescription for this unit was to remove all merchantable stems except 3 mature white birch per ha, 1-2 mature red oak per ha, and some healthy beech (~7 per ha). Mature white birch in this unit were in the latter stages of decline before harvesting, but those retained were intended to act as a seed source during the post-harvest period. Retained beech show little to no

sign of bark disease and may represent resistant genotypes. Retained red oak were intended as a seed source, especially important in light of the loss of most beech to bark disease. Cutting on this unit was completed in October-November 2019.

Units 1, 5, 11, 14, 15, 16, 17 Concept: *Over-mature Reference*. These units include the full range of ecosystem and cover types, landforms, soil and hydrologic conditions present on the units to be treated, and afford the opportunity to continue monitoring aspects of these ecosystems as they continue to advance through succession, which will increasingly involve the blowdown and breakup of over-mature aspen. Future management activities could be conducted in areas of units 15 or 17, but if so, that management should be restricted to areas that have no long-term monitoring locations (e.g., plots).

Units 2, 4, 6, 8, 10, 12, 14, 18, 19 Concept: *Young Reference*. These units include the full range of ecosystem types, landforms, soil and hydrologic conditions present on the units to be treated, and afford the opportunity to continue monitoring aspects of these ecosystems as they begin to enter mid-successional stage. Because these are younger stands (cut in the 1970s and 1980s), they may be useful as a “time machine” of sorts, either in comparison to the over-mature reference units, or to the treatment units harvested in 2019. Given the good growth rates of aspen currently dominating these units, it is conceivable that merchantable biomass could be harvested from them within the next 10 years, especially if included in a larger timber sale involving other units.

Overall, this set of proposed treatments will provide immediate opportunities to study management impacts on hydrology, biogeochemistry, and organismal behavior in forest ecosystems. During the years following treatments, additional treatments may be considered for portions of the management area not presently proposed for manipulation. If realizing the significant revenue currently held in the mature aspen is a management goal, additional harvesting should be conducted soon, since these stands are beginning to break up and the large trees are susceptible to windthrow. Regardless whether additional revenue is desired, the landscape mosaic of stands in the overall management area will create more heterogeneous forest conditions, to the benefit of the landscape and its researchers. Below is an overview research questions and activities that have been conducted or are planned.

Research questions: These treatments integrate with ongoing and create new opportunities for research projects. The project team will continue publishing papers on soil and ecosystem hydrology and biogeochemistry. Management treatments will address objectives of the original McIntire-Stennis grant (Drevnick, Kerkez, Nave) that funded the initiation of the HCW study, pertaining to the effects of harvesting on the hydrologic movement of C and Hg. Key projects completed or ongoing are:

1. Forest inventory: The 2018 AAME team (Olivia Brinks, John Den Uyl, Elizabeth Feliciano, Edie Juno, Katy Hofmeister, Luke Nave, and Adam Schubel) marked and GPS'd 53 permanent plots (10 m radius each) distributed across the management units (blue points in Fig. 2). Data recorded includes: 1) for every standing tree >8 cm dbh: species, dbh, status (live or dead), azimuth from plot center, approximate distance, whether marked for retention during harvest, and if beech, the density of scale infestation [3 categories]; 2) for every tree <8 cm dbh but reaching dbh, the species and size class (0-2, 2-4, 4-6, or 6-8 cm dbh). These data were the basis

for Elizabeth Feliciano's Frontiers project at UMBS, and provide a complete forest inventory baseline before management treatments occurred. Data are available upon request by any researcher.

2. Soil sampling for post-hoc experiment: The 2018 AAME team sampled 25 soil profiles (to E'Bt horizon or 2 m depth, whichever is shallower) from Units 2, 3, 4, 15, and 16. This sample set is being used to test for impacts of past harvesting on the storage and stability of C in multiple soil C pools. Laboratory processing has been completed and extractions and analyses are in progress. Analyses for this study are expected to run into 2020; most will be performed in the UMBS Analytical Lab, and some will occur through NIACS' Carbon, Water, and Soils Lab. This will eventually yield a novel paper on multiple mechanisms of soil C stabilization in Spodosols and accompanying analysis of management effects on soil C stability.
3. Groundwater, porewater, and soil sampling for real-time experiment: Den Uyl, Hofmeister, and Nave augmented existing monitoring locations in Units 12 (1980s harvest), 13 (2019 harvest), and 14 (reference) for a study of the real-time effects of management on water table dynamics, soil C stability and transport in pore- and groundwater. In these units, 13 water table wells are installed to the restrictive layer (till) or to 2m (whichever is shallower); 12 possess water level loggers. Soils have been collected, processed and archived for all of these locations. Tension lysimeters were deployed to the B1 horizon (~30 cm) at 12 of the locations in fall 2018 by Den Uyl. In 2019, Technician Brooke Propson, Adam Schubel, and Nave deployed 3 new wells on private property ~1km south of Honeysuckle Creek. This property, owned by Lynn Lord (summer resident of W. Burt Lake Rd.) and overseen by UMBS friend /year-round resident Rick Johnson, was logged in 2018. Soils were sampled while installing these wells, which are paired with the three along Honeysuckle Creek by elevation and landscape ecosystem type.
4. Groundwater and porewater sampling: Biweekly to monthly collections of groundwater and porewater (from wells and lysimeters) are planned to begin with spring melt 2020. These water samples, along with the soils from the water sampling locations, will be analyzed for C forms and fluxes for a paper documenting the real-time changes in C hydrobiogeochemistry following harvest. As of this writing, 4 collections have been made (2x in May 2019, 1x each in October and November 2019).
5. Microclimate sampling: Brooke Propson, John Den Uyl, and Nave collected soil moisture (May 2019) and soil moisture, temperature, air temperature, humidity, and sunlight data (July 2019) across 53 permanent plots. Data are planned to be collected at more regular intervals in 2020 beginning with snowmelt. This plot-based distributed data collection is to augment, and capture spatial variability in meteorological parameters being recorded at half-hourly intervals by weather / soil stations in Units 12, 13, 14, which were installed in Autumn 2019. The stations are deployed at similar elevation, slope and aspect, allowing for comparison of real-time weather variables in the Resistance cut, ~35 year old, and reference aspen areas.
6. Coarse woody debris: Collecting woody debris data is a priority for 2020; this is planned to be done by the Forest Ecosystems class using the new permanent sample plots where trees have been inventoried.

References and additional resources:

NIACS' Forest Adaptation Workbook (<https://adaptationworkbook.org/>). The Adaptation Workbook is a web-based process for incorporating climate change adaptation into forest planning. I have used the Adaptation Workbook as I have developed this management plan (with special help from Stephen Handler, Maria Janowiak, Dean Reid and Adam Schubel) and will share it / walk through it with anyone interested.

Handler S, Duveneck MJ, Iverson L, Peters E, Scheller RM, Wythers KR, Brandt L, Butler P, Janowiak M, Shannon PD, Swanston C, Eagle AC, Cohen JG, Corner R, Reich PB, Baker T, Chhin S, Clark E, Fehring D, Fosgitt J, Gries J, Hall C, Hall KR, Heyd R, Hoving CL, Ibáñez I, Kuhr D, Matthews S, Muladore J, Nadelhoffer K, Neumann D, Peters M, Prasad A, Sands M, Swaty R, Wonch L, Daley J, Davenport M, Emery MR, Johnson G, Johnson L, Neitzel D, Rissman A, Rittenhouse C, Ziel R. 2014. *Michigan forest ecosystem vulnerability assessment and synthesis: a report from the Northwoods Climate Change Response Framework project*. Gen. Tech. Rep. NRS-129. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 229 pp. <https://www.nrs.fs.fed.us/pubs/45688>

Hofmeister K.L., Nave L.E., Riha S.J., Schneider R.L., Walter M.T. 2019. A Test of Two Spatial Frameworks for Representing Spatial Patterns of Wetness in a Glacial Drift Watershed. *Vadose Zone Journal* 18: 180054

Hofmeister K.L., Nave L.E., Drevnick P.E., Veverica T.J., Knudstrup R.V., Heckman K.A., Riha S.J., Schneider R.L., Walter M.T. 2019. Seasonal dynamics and exports of elements from a first-order stream to a large inland lake in Michigan. *Hydrological Processes* 33: 1476-1491

Nave L.E., Gough C.M., Perry C.H., Hofmeister K.L., Le Moine J., Domke G.M., Swanston, C.W., Nadelhoffer K.J. 2017. *Physiographic factors underlie rates of biomass production during succession in Great Lakes forest landscapes*. *Forest Ecology and Management* 397: 157-173

Pearsall D.R. 1995. *Landscape ecosystems of the University of Michigan Biological Station: ecosystem diversity and ground –cover diversity*. School of Natural Resources and Environment, University of Michigan, Ann Arbor. Dissertation, 396pp., plus maps.

Swanston CW, Janowiak MK, Brandt LA, Butler PR, Handler SD, Shannon PD, Lewis AD, Hall K, Fahey RT, Scott L, Kerber A, Miesbauer JW, Darling L. 2016. *Forest Adaptation Resources: Climate Change Tools and Approaches for Land Managers, 2nd edition*. General Technical Report NRS-87-2. U.S. Department of Agriculture, Forest Service, Newtown Square, PA. 170pp. <https://www.nrs.fs.fed.us/pubs/40543>